Binge Drinking and Other Risk Behaviors among College Students

2014

April 2014

Prepared by the University of Delaware Center for Drug & Alcohol Studies

Key Staff for the 2014 College Risk Behaviors Study

(In alphabetical order)

Roberta Gealt
Whitney Gunter
Steven Martin
Daniel O'Connell
Christy Visher

Center for Drug and Alcohol Studies
University of Delaware
Newark, DE 19716-2582
(Address all inquiries to Steven S. Martin)

This research was sponsored by the Substance Abuse and Mental Health Services Administration Center for Substance Abuse Prevention. The views and conclusions expressed in this report do not necessarily represent those of the sponsoring agency.

College Risk Behaviors Study

Introduction

The College Risk Behaviors Study (CRBS) was administered during the 2014 spring semester to a random sample of full-time, undergraduate, University of Delaware students. Each was asked via email to follow an enclosed link and complete the online survey. These students were offered \$5

in flex credit as compensation for their time. By the end of the semester, 1,121 students chose to participate, corresponding to a 37% response rate, a rate equivalent to similar online surveys. Given the chance to specifically opt out of the survey, only 1% of students chose to do so.

The sample is similar to the overall student body in distribution of gender and race/ethnicity, with slightly higher percentages of females and white students in the sample than in the total UD student population. It also includes a somewhat greater percentage of students living on-campus than does the population. Due to the large difference in gender, the data were weighted to adjust the sample gender ratio to match that of the population's.


This report will address several issues relating to substance consumption using data from these participants. First, consumption prevelance rates will be presented for varying levels of alcohol consumption, with gender and race/ethicity differences and similarities highlighted. Second, an indepth analysis of the potential consequences of alcohol use and abuse will illustrate the negative impact of these behaviors. Finally, data from past studies will be presented to indicate trends trends over the past 20 years at University of Delaware.


survey. These students were offered		
Characteristics of the Sample*		
	Sample	UD
Gender		
Male	34%	42%
Female	66%	58%
Race		
White	77%	77%
Black	4%	5%
Hispanic	7%	6%
Asian	9%	4%
Other	3%	8%
Residence		
On-Campus	52%	44%
Off-Campus	48%	56%
Class Year		
Freshmen	21%	21%
Sophomore	24%	23%
Junior	27%	30%
Senior	27%	26%
Age		
18	13%	
19	25%	
20	27%	
21	23%	
22	10%	
23 or Older	3%	

^{*} Aside from gender, enrollment statistics for demographics in 2013-2014 were not yet available. Statistics on race, residency, class year, and age for the university are from 2012-2013.


Alcohol Consumption Prevalence Rates

In general, the substance use of choice among students was alcohol. Though approximately 19% of students reported using illegal drugs, the vast majority were also using alcohol, with less than 1% of students using just illegal drugs. Of students using drugs, about three-quarters were using only marijuana.






Alcohol was the most used substance, with 78% of students reporting drinking alcohol in the past month. The majority of students (61%) binge drank five or more drinks in a single sitting during the past month. Males (65%) were significantly more likely to engage in binge drinking than were females (58%). Heavy alcohol use (drinking on at least two-thirds of the past 30 days) is much lower, with 3% of students reporting heavy drinking habits.





Additional differences emerge when alcohol use is examined by race/ethnicity. Past month alcohol consumption was much more common among non-Hispanic Whites (84%) than it was for minority students (60%). This is also true of binge drinking behaviors, with approximately two-thirds (68%) of White students reported binge drinking, and slightly less than one-third (38%) of students identifying with other races reported similar drinking.



Non-Hispanic White students are more likely to drink or binge drink than other ethnicities.


When broken down by class year, a steady increase in alcohol consumption can be seen over time. This applies to drinking in general, binge drinking and, to a lesser extent, heavy alcohol use.





Binge drinking appears steady throughout the college experience. However, casual drinking and heavy drinking tend to increase over time.

^{*} Class difference is significant at the .05 level




Casual drinking, binge drinking and heavy drinking tend to increase as students get older.

^{*} Race difference is significant at the .05 level

 $^{^{}st}$ Age difference is significant at the .05 level

Students report getting alcohol from various sources in various ways.* For those underage, parties and friends are the most likely sources, followed closely by having someone else (presumably also a friend) buy it for them. Turning 21 opens legal purchasing options at restaurants, bars and liquor stores. Those age 21 and older are much more likely to buy from these sources, and relatedly less likely to rely on parties and friends. These informal sources, however, still remain a source among these older students.





Also worth noting is that the use of false identification to obtain alcohol is low. When looking only at those too young to purchase alcohol legally, only 5% of students report using a fake ID to purchase the alcohol at a restaurant, bar or liquor store.†

^{*} Students were asked, hypothetically, if they wanted alcohol, where they would get it. The percentages reported here reflect the proportion of alcohol-consuming students who reported they would "often" or "very often" get alcohol from these sources.

Consequences of Alcohol Consumption

The College Risk Behaviors Study also measured several potential consequences of alcohol consumption and binge drinking. The data below present a comparison between binge drinkers and non-binge drinkers for the likelihood of experiencing these consequences. Binge drinking was defined as drinking five or more drinks in a single sitting. Occasional binge drinkers include students who binge drank one to nine times in the past month, while students who binge drank more often were considered frequent binge drinkers.






^{*} Victimization questions were defined as the following – Assault: Being attacked or someone threatening to attack you (11%) – Sexual Assault: Having been forced or coerced to engage in unwanted sexual intercourse of any kind including vaginal, anal, or oral sex (6%). All were limited to incidents occurring in the past year.

Consumption Trends

The College Risk Behaviors Study was not the first survey of University of Delaware students. In 1993, from 1997 to 2003, and again in 2005, University of Delaware participated in the College Alcohol Study. As the trends below indicate, cigarette use has steadily declined over time.




Binge drinking saw a decline in the late 1990s and early 2000s, but has returned to its previous rate. Cigarettes use experienced a steady decline, as has marijuana use.

Note: Data from 1993, 1997-2003, and 2005 are from the College Alcohol Study. Data beginning in 2008 are from the College Risk Behaviors Study.

Previous reports used unweighted data. For these trend lines, the data from 2011 to the present were adjusted to include weights for gender.

Compared to national trend data, University of Delaware binge drinking rates are consistently higher than the national average. Even during the decline in the late 1990s, UD's rate was significantly higher than rates calculated nationally. The exact reason for this is not clear and it may be a combination of several factors. It is possible, for example, that other institutions in the north-east area with a high proportion of students living on-campus are also above the national average to some extent. Unfortunately, data that separate out institutions similar to University of Delaware are not currently publicly available.



The changes in binge drinking rates at University of Delaware appear to be a local phenomenon and not something occurring nationally.

Note: National data from the College Alcohol Study (CAS) and from the Monitoring the Future (MTF), 1993-2011.

The timeframe for binge drinking is defined differently between studies. The UD (1993-2005) and CAS surveys used a timeframe of past two weeks, while the UD (2008-Present) and MTF surveys used past month. Though the timeframe varies, there does not appear to be any increase for using past month rather than past two weeks based on similar numbers from the CAS and MTF.